

Cardiovascular Consequences of Obstructive Sleep Apnea (Current Topics in Sleep Medicine from the University of Pennsylvania)  
Samuel T. Kuna, MD  
References

### Chapter 1

1. Motta J, Guilleminault C, Schroeder JS, Dement WC. Tracheostomy and Hemodynamic Changes in Sleep-Induced Apnea. *Annals of Internal Medicine*. 1978;89(4):454-8.
2. Mehra R, Benjamin EJ, Shahar E, Gottlieb DJ, Nawabit R, Kirchner HL, et al. Association of nocturnal arrhythmias with sleep-disordered breathing: The Sleep Heart Health Study. *Am J Respir Crit Care Med*. 2006 Apr 15;173(8):910-6.

### Chapter 2

1. Gami AS, Howard DE, Olson EJ, Somers VK. Day–Night Pattern of Sudden Death in Obstructive Sleep Apnea. *New England Journal of Medicine*. 2005;352(12):1206-14.
2. Young T, Finn L, Peppard PE, Szklo-Coxe M, Austin D, Nieto FJ, et al. Sleep disordered breathing and mortality: eighteen-year follow-up of the Wisconsin sleep cohort. *Sleep*. 2008 Aug;31(8):1071-8.
3. Redline S, Yenokyan G, Gottlieb DJ, Shahar E, O'Connor GT, Resnick HE, et al. Obstructive Sleep Apnea–Hypopnea and Incident Stroke. *American Journal of Respiratory and Critical Care Medicine*. 2010 /07/15;182(2):269-77.
4. Gottlieb DJ, Yenokyan G, Newman AB, O'Connor GT, Punjabi NM, Quan SF, et al. Prospective study of obstructive sleep apnea and incident coronary heart disease and heart failure: the sleep heart health study. *Circulation*. 2010 Jul 27;122(4):352-60.
5. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet*. 2005 Mar 19-25;365(9464):1046-53.

### Chapter 3

1. Savransky V, Nanayakkara A, Li J, Bevans S, Smith PL, Rodriguez A, et al. Chronic intermittent hypoxia induces atherosclerosis. *Am J Respir Crit Care Med*. 2007 Jun 15;175(12):1290-7.
2. Schulz R, Murzabekova G, Egemenazarov B, et al. Arterial hypertension in a murine model of sleep apnea: role of NADPH oxidase 2. *J Hypertens*. 2014 Feb;32(2):300-5.
3. Hedner J, Darpo B, Ejnell H, Carlson J, Caidahl K. Reduction in sympathetic activity after long-term CPAP treatment in sleep apnoea: cardiovascular implications. *European Respiratory Journal*. 1995 February 1, 1995;8(2):222-9.

## Chapter 4

1. Nieto F, Young TB, Lind BK, et al. Association of sleep-disordered breathing, sleep apnea, and hypertension in a large community-based study. *JAMA*. 2000;283(14):1829-36.
2. Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *New England Journal of Medicine*. 2000;342(19):1378-84.
3. Marin JM, Agusti A, Villar I, et al. Association between treated and untreated obstructive sleep apnea and risk of hypertension. *JAMA*. 2012;307(20):2169-76.
4. Pepperell JCT, Ramdassingh-Dow S, Crosthwaite N, Mullins R, Jenkinson C, Stradling JR, et al. Ambulatory blood pressure after therapeutic and subtherapeutic nasal continuous positive airway pressure for obstructive sleep apnoea: a randomised parallel trial. *Lancet*. 2002;359(9302):204-10.
5. Fava C, Dorigoni S, Vedove FD et al. Effect of CPAP on blood pressure in patients with OSA/hypopnea: A systematic review and meta-analysis. *Chest* 2014;145(4):762–771
6. Iftikhar IH, Valentine CW, Bittencourt LR, et al. Effects of continuous positive airway pressure on blood pressure in patients with resistant hypertension and obstructive sleep apnea: a meta-analysis. *J Hypertens*. 2014 Dec;32(12):2341-50.
7. Chirinos JA, Gurubhagavatula I, Teff K, et al. CPAP, weight loss, or both for obstructive sleep apnea. *N Engl J Med*. 2014 Jun 12;370(24):2265-75.