

## Chapter 2

1. Boeve BF. REM sleep behavior disorder: Updated review of the core features, the REM sleep behavior disorder-neurodegenerative disease association, evolving concepts, controversies, and future directions. *Ann N Y Acad Sci.* 2010 Jan;1184:15-54.
2. Iranzo A, Santamaria J. Severe obstructive sleep apnea/hypopnea mimicking REM sleep behavior disorder. *Sleep.* 2005 Feb;28(2):203-6.

## Chapter 3

1. Boeve BF, Silber MH, Saper CB, Ferman TJ, Dickson DW, Parisi JE, et al. Pathophysiology of REM sleep behaviour disorder and relevance to neurodegenerative disease. *Brain.* 2007 Nov;130(Pt 11):2770-88.

## Chapter 4

1. Boeve BF, Silber MH, Saper CB, Ferman TJ, Dickson DW, Parisi JE, et al. Pathophysiology of REM sleep behaviour disorder and relevance to neurodegenerative disease. *Brain.* 2007 Nov;130(Pt 11):2770-88.
2. Schenck CH, Bundlie SR, Mahowald MW. Delayed emergence of a parkinsonian disorder in 38% of 29 older men initially diagnosed with idiopathic rapid eye movement sleep behaviour disorder. *Neurology.* 1996 Feb;46(2):388-93.
3. Schenck CH, Boeve BF, Mahowald MW. Delayed emergence of a parkinsonian disorder or dementia in 81% of older men initially diagnosed with idiopathic rapid eye movement sleep behavior disorder: a 16-year update on a previously reported series. *Sleep Med.* 2013 Aug;14(8):744-8.
4. Boeve BF, Silber MH, Ferman TJ, Lucas JA, Parisi JE. Association of REM sleep behavior disorder and neurodegenerative disease may reflect an underlying synucleinopathy. *Mov Disord.* 2001 Jul;16(4):622-30.
5. Postuma RB, Gagnon JF, Vendette M, Montplaisir JY. Markers of neurodegeneration in idiopathic rapid eye movement sleep behaviour disorder and Parkinson's disease. *Brain.* 2009 Dec;132(Pt 12):3298-307.
6. Iranzo A, Fernandez-Arcos A, Tolosa E, Serradell M, Molinuevo JL, Valldeoriola F, et al. Neurodegenerative disorder risk in idiopathic REM sleep behavior disorder: study in 174 patients. *PLoS One.* 2014;9(2):e89741.
7. Iranzo A, Lomena F, Stockner H, Valldeoriola F, Vilaseca I, Salamero M, et al. Decreased striatal dopamine transporter uptake and substantia nigra hyperechogenicity as risk markers of synucleinopathy in patients with idiopathic rapid-eye-movement sleep behaviour disorder: a prospective study [corrected]. *Lancet Neurol.* 2010 Nov;9(11):1070-7.
8. Holtbernd F, Gagnon JF, Postuma RB, Ma Y, Tang CC, Feigin A, et al. Abnormal metabolic network activity in REM sleep behavior disorder. *Neurology.* 2014 Feb 18;82(7):620-7.
9. Postuma RB, Gagnon JF, Rompre S, Montplaisir JY. Severity of REM atonia loss in idiopathic REM sleep behavior disorder predicts Parkinson disease. *Neurology.* 2010 Jan 19;74(3):239-44.

10. Fantini ML, Gagnon JF, Petit D, Rompre S, Decary A, Carrier J, et al. Slowing of electroencephalogram in rapid eye movement sleep behavior disorder. *Ann Neurol*. 2003 Jun;53(6):774-80.
11. Martín RU-S, Francke PV, Illanes FL, Gazmuri AJ, Rivera JS, Ferrández JG, et al. Plasma urate in REM sleep behavior disorder. *Movement Disorders*. 2013;28(8):1150-1.

## Chapter 5

1. McCarter SJ, St Louis EK, Boswell CL, Dueffert LG, Slocumb N, Boeve BF, et al. Factors associated with injury in REM sleep behavior disorder. *Sleep Med*. 2014 Nov;15(11):1332-8.
2. Standards of Practice Committee, R. Nisha Aurora, Rochelle S. Zak, Rama K. Maganti, Sanford H. Auerbach, Kenneth R. Casey, Susmita Chowdhuri, Anoop Karippot, Kannan Ramar, David A. Kristo, Timothy I. Morgenthaler. Best Practice Guide for the Treatment of REM Sleep Behavior Disorder (RBD). *J Clin Sleep Med*. 2010 February 15; 6(1): 85–95.